

The association between lower urinary tract symptoms and falls: forming a theoretical model for a research agenda

Gibson, William; Hunter, Kathleen F. ; Camicioli, Richard; Booth, Joanne; Skelton, Dawn A.; Dumoulin, Chantale; Paul, Lorna; Wagg, Adrian

Published in:
Neurourology and Urodynamics

DOI:
[10.1002/nau.23295](https://doi.org/10.1002/nau.23295)

Publication date:
2018

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Gibson, W, Hunter, KF, Camicioli, R, Booth, J, Skelton, DA, Dumoulin, C, Paul, L & Wagg, A 2018, 'The association between lower urinary tract symptoms and falls: forming a theoretical model for a research agenda', *Neurourology and Urodynamics*, vol. 37, no. 1, pp. 501-509. <https://doi.org/10.1002/nau.23295>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

THE ASSOCIATION BETWEEN LOWER URINARY TRACT SYMPTOMS AND FALLS: FORMING A THEORETICAL MODEL FOR A RESEARCH AGENDA

William Gibson¹, Kathleen F. Hunter², Richard Camicioli³, Joanne Booth⁴, Dawn A. Skelton⁴, Chantale Dumoulin⁵, Lorna Paul⁴, Adrian Wagg¹

¹ Division of Geriatric Medicine, Department of Medicine, University of Alberta, Edmonton, Alberta, Canada

² Faculty of Nursing, University of Alberta, Edmonton, Alberta, Canada

³ Division of Neurology, University of Alberta, Edmonton, Alberta, Canada

⁴ School of Health & Life Sciences, Glasgow Caledonian University, Glasgow, UK

⁵ Faculty of Medicine, Université de Montréal, Montréal, Canada

Corresponding Author:

wgibson@ualberta.ca, +1 780 492 0133

THE ASSOCIATION BETWEEN LOWER URINARY TRACT SYMPTOMS AND FALLS: FORMING A THEORETICAL MODEL FOR A RESEARCH AGENDA

ABSTRACT

There is a well-recognised association between falls and lower urinary tract symptoms (LUTS) in older adults, with estimates of odd ratios for falls in the presence of LUTS ranging between 1.5 and 2.3. Falls and LUTS are both highly prevalent among older people and are both markers of frailty, with significant associated morbidity, mortality, and healthcare resource cost. Despite the recognition of this association, there has been little research to examine the potential causes of this association, and no intervention trial has established if reducing LUTS or urinary incontinence can reduce the risk of falls. In this paper, stemming from a meeting of experts in the field, the current evidence base and hypothesized causal linkage for this association is reviewed and an outline for a potential research agenda to advance the science in the area is forwarded.

INTRODUCTION

Urinary incontinence (UI) and lower urinary tract symptoms (LUTS), including urinary urgency, frequency, and nocturia are highly prevalent amongst the general population; this prevalence rises in association with increasing age^{1,2}. LUTS and UI are stigmatizing conditions³, which are often under-reported and under-treated, particularly in older individuals^{4,5}. The most common form of UI in older people is urgency urinary incontinence (UUI), urine loss associated with urinary urgency; a sudden, overwhelming desire to void that is difficult to defer⁶. Frequency, urgency and nocturia, the most common LUTS, which are also components of overactive bladder syndrome (OAB)⁶, are extremely common in later life; with up to 50% of men and 60% of women aged over 70 years old describing at least one lower urinary tract symptom¹. The diagnosis of OAB requires the individual to report urinary urgency.

Up to one third of people aged over 65 years, and half of those over 80 years old, will fall in any given year⁷. Falls, defined by the World Health Organisation as “an event which results in a person coming to rest inadvertently on the ground or floor or other lower level”⁸ are often recurrent, with around half of people who fall experiencing another within 12 months⁹. Falls negatively affect quality of life, cause individual pain and suffering, lead to functional decline¹⁰, and are associated with significant healthcare resource use. In the UK, 38% of the 816 participants in the Newcastle 85+ Cohort Study had fallen in the previous year, with 10% of those having fallen suffering a fracture at an average cost of £200 per person (2007 prices)¹¹. In the USA, there were approximately 24,190 fatal and 3.2 million non-fatal falls among older adults, at a total cost of over US\$30billion in 2105¹². In Canada it is predicted that by 2031, the annual direct healthcare costs attributable to falls will be C\$4.4 billion¹³, and in 2004, falls were responsible for 7.3% of all hospital admissions for older people and the annual direct cost of fall-related injuries was estimated at C\$2billion¹³.

There is a well-recognised association between falls and lower urinary tract symptoms in older adults¹⁴⁻²¹. Older people with urinary urgency or UUI are significantly more likely to fall and sustain injury compared to age-matched controls, with estimates of the odds ratio for falls ranging from 1.5 to 2.3^{15,22,23}. However, the reasons for this association are neither understood nor well-studied. This paper is the result of a meeting of health professionals and researchers with expertise in LUTS in older people, exercise physiology, cognitive processing, rehabilitation and bladder physiology. Here,

the current state of the research evidence is reviewed, a theoretical framework to explain the association is proposed, and a research agenda to better explain this phenomenon outlined.

WHAT IS KNOWN ABOUT THE ASSOCIATION BETWEEN FALLS AND LUTS?

Brown and colleagues¹⁴ performed a secondary analysis of data from an osteoporosis cohort study, examining a group of 6049 community-dwelling older women using regular self-completed questionnaires sent to all participants every four months. In this cohort, followed for an average of three years, those with at least one weekly urgency urinary incontinence (UUI) episode were more likely to fall (OR 1.26, 95%CI 1.14-1.40) than those without. Weekly UUI was also associated with higher odds of sustaining a non-spinal fracture (hazard ratio 1.34 95%CI 1.06-1.69). In this study, stress incontinence was not associated with higher odds of falling (Odds ratio (OR) 1.06, CI 0.95-1.19) or sustaining a fracture (relative hazard 0.98, CI 0.75-1.28). Analysis of the Concord Health and Ageing in Men Project, a longitudinal study of community-dwelling men in Australia followed 1090 men over a period of 2 years. Here, the presence of urgency incontinence, defined as weekly episodes of UUI, was associated with a higher incidence of falls (OR 2.57 95%CI 1.51 – 4.3) and men with a higher International Prostate Symptom Score storage sub-score, defined a score of 19 and above, had a higher incident rate of falls (incident rate ratio 1.72 (95%CI 1.24-2.38)²⁴. A Japanese study of patients with Parkinson's disease found that increased micturition frequency either by day or night was not associated with falls, but that the presence of urinary urgency was strongly associated with a large increase in the odds of falling (OR 5.14 95%CI 1.51-17.48). Only 14% of the falls reported in this study occurred on the way to or from the toilet²⁵. There appears to be no association between voiding LUTS and falls in men based upon current published evidence.

A recent systematic review of the association between falls and LUTS in community-dwelling men aged 60 years and over identified six cross-sectional studies and three prospective cohort studies. The identified data were only suitable for qualitative synthesis but urinary incontinence and storage LUTS were consistently shown to have a weak to moderate association with an increased likelihood of falls. None of the identified studies examined potential causes for these associations; the categorisation of continence or not and degree of accounting for confounding variables was inconsistent across the included studies²⁶. A small cross-sectional analysis of community dwelling women aged 65 and over in the US examined the association between nocturia, nocturnal enuresis and falls. Neither severity of UI nor severity of nocturia was associated with an increased risk of falls, but there was a statistically significant association between nocturnal enuresis and impairment of physical function and the presence of frailty. However, in the multivariable regression model, which included age, physical function, and the frequency of nocturnal enuresis episodes, only physical function remained as significant risk factor for falls²⁷.

The use of sedative medications has been identified as a potential risk factor for falls in those with LUTS. In long-term users of benzodiazepines, urinary incontinence and LUTS were an independent risk factor for falls in older adults irrespective of the degree of exposure to these drugs. An exposure-response relationship was observed between the frequency of incontinence and falls, with falls occurring in 17% of participants with incontinence only once weekly, 25% of those with incontinence two to three times per week, 60% of those with incontinence daily, and 58% of those with incontinence more than once per day²⁸.

There is, therefore, evidence for a reasonably strong association between falls and LUTS/UI in older adults, but the mechanism underlying this is unexplained. Little research has addressed this and additionally, despite calls for such work to be undertaken, no intervention trial has yet been performed to examine if treating LUTS can reduce the risk of falls^{29,30}.

POTENTIAL LINKS AND AVENUES FOR FURTHER STUDY

RUSHING TO THE TOILET OR SLIPPING IN URINE

The idea that the reason that people with incontinence are more likely to fall is due to rushing to get to the toilet or by slipping in their own urine following an episode of incontinence has been cited in previous publications as an explanation of the association between LUTS and falls^{15,31} without either evidence or clarification. These were reviews, citing no data, and thus there is no evidence in the published literature to suggest that this is the case.

A case-control study in the UK found no temporal association between getting to the toilet and falling, suggesting that a simplistic explanation such as rushing is unlikely to be the underlying link between the two entities³². In this study only 6% of participants identified a temporal relationship between their fall and experience of urgency. This is similar to the finding of Sakushima et al²⁵, which reported 14% of falls related to getting to or from the toilet, but did not examine whether or not those people perceived themselves to be "rushing". In addition, there is evidence from continent, middle aged women that the response to a strong desire to void is to slow down, not to speed up or rush³³. The idea that "rushing" explains the association between urgency and falls should therefore be viewed with caution until evidence supporting this intuitive response is produced. , At present, there is no evidence to support rushing as a link between urgency and falls, and the limited evidence available suggests that rushing is not implicated in the association.

ACTIVITY RESTRICTION

Lower levels of physical activity have been associated with increased falls³⁴ and there is evidence that fear of falling is associated with decreased activity in older adults^{34,35}. For some people there is consequent avoidance of activity resulting from the fear of falling³⁶, which in turn further impairs physical ability and increases the risk of future falls³⁷. Women with incontinence will often limit their participation in physical activity and sport as a result of their incontinence³⁸, and recent work found that older women with UI had worse performance on tests of balance and reported more falls than continent, age-matched controls³⁹. There is evidence that the presence of urinary incontinence is associated with limitations in activities of daily living⁴⁰, and that incontinence in older adults is associated with deconditioning, the development of obesity and a decline in general health⁴¹. However, some studies have only found reduced levels of physical activity associated with stress, but not urgency incontinence⁴². For example, a French cohort study of 1942 community-dwelling women aged between 75 and 85 used a panel of balance assessments and the International Consultation on Incontinence Questionnaire–Short Form questionnaire (ICIQ-SF) to assess the associations between physical limitation and LUTS. The authors reported a significant deterioration in all the standard mobility and balance test results according to the severity of UI, including timed up and go, standing balance, walking balance, and single leg stand time⁴³.

Given that deconditioning and reduced physical fitness is associated with an increase in falls risk⁴⁴, responding to urinary urgency in a deconditioned older person may, in itself, increase the risk of a subsequent fall. Exercise interventions have been demonstrated to be effective in reducing the

frequency of incontinence episodes in nursing home residents⁴⁵ and community-dwelling older women⁴⁶. A short-term (8-week) intervention of physiotherapy-delivered training in mobility and toileting skills in non-demented older women living in a care facility resulted in a 37% reduction in the daily urine loss⁴⁷. Likewise there is good evidence that increasing physical activity can decrease falls in older adults living in the community⁴⁸, in institutions⁴⁹ and in individuals with dementia⁵⁰. A systematic review of cost-effectiveness of falls studies concluded that physical activity, specifically the Otago Exercise programme in those 80 years and older, was one of three cost saving falls prevention strategies⁵¹.

Current evidence tentatively supports a bi-directional relationship between urinary incontinence and activity restriction, with those experiencing incontinence reducing their trips outside the home and those with limited mobility experiencing more incontinence.

FRAILITY, MULTIMORBIDITY, AND POLYPHARMACY

It is undeniable that urinary incontinence, and in particular urgency incontinence, can be considered as a geriatric syndrome and a marker of frailty⁵² and, furthermore, frailty is strongly associated with falls^{53,54}. Frailty, defined by Fried as “a biologic syndrome of decreased reserve and resistance to stressors, resulting from cumulative declines across multiple physiological systems, and causing vulnerability to adverse outcomes”⁵⁵ is not synonymous with multimorbidity, or disability, but rather describes a state of increased vulnerability to adverse outcome. Many tools exist for quantifying and diagnosing frailty⁵⁶⁻⁵⁸, a detailed discussion of which is beyond the scope of this review.

It is conceivable that no causal link exists between falls and incontinence or LUTS, and that these are both markers of frailty. UI and UUI have been associated with musculoskeletal conditions including low back pain and osteoarthritis in a cross-sectional study of older women in Japan⁵⁹. A retrospective cross sectional study of older patients admitted to hospital in Germany with a fracture found that those with UI were frailer, more dependent, and had higher levels of physical and cognitive impairment than those without UI⁶⁰. Both frailty and LUTS are strongly associated with multimorbidity and polypharmacy^{55,61,62}, which in turn are influenced by the presence of vascular risk factors⁶³ and inflammation⁶⁴. A retrospective analysis of drug dispensing data in a Japanese cohort found that polypharmacy was associated with the use of medications known to contribute to urgency⁶⁵. Similarly, a Canadian cross-sectional study identified a strong association (OR 4.9, 95%CI 3.1-7.9) between polypharmacy, defined as 5 or more medications, and the prescription of a medication known to cause LUTS⁶⁶. However, there is little evidence that polypharmacy in the absence of drugs that cause incontinence has an impact on the lower urinary tract.

Several other specific comorbidities are associated with both LUTS and falls. For example, as well as being associated with arterial disease, diabetes mellitus has well documented effects on the lower urinary tract and is associated with impaired bladder emptying and urgency⁶⁷, and can cause neuropathies that may predispose to falls including peripheral neuropathy causing gait disturbance and autonomic neuropathy causing postural hypotension⁶⁸. Likewise, Parkinson’s disease is commonly associated with both LUTS⁶⁹ and falls⁷⁰.

These relationships are summarised in figure 1. The observed association between falls and LUTS in older adults may therefore be due to the confounding effect of their multiple common causes. However, given that the epidemiological evidence for this association is overwhelmingly from

community-based studies, the majority of available data come from non-frail individuals. Brown et al⁷¹ identified 20% of their cohort as being frail (defined by the investigators based being “weak, unsteady, and fragile”) at baseline, and Noguchi et al, although not specifically identifying frailty as a measure at enrolment, described a cohort in which 97% of participants were mobile without the use of a walking aid²⁴. A case-control study of nursing home residents in the USA found that those with UI or OAB exhibited higher rates of cognitive impairment, mobility impairment, and higher numbers of comorbidities than those without⁷². It is, therefore, reasonable to suggest that although frailty is a common cause of both falls and LUTS/UI, there are other complex contributors to the association, including the role of cognitive influences and potentially other, as yet unknown, links. The paucity of evidence from frail or institutionalised older people limits the applicability of the current evidence base to this population.

CENTRAL CONTROL AND COGNITIVE FACTORS

The maintenance of continence is not an innate ability; much like walking, it is a learned skill developed in the early years of life, and once attained, continence is under conscious control⁷³. Numerous areas of the brain are involved in the maintenance of continence, with research techniques including brain imaging, functional brain imaging, and evidence from disease-specific studies being used to delineate the complex relationships between them⁷⁴. Functional positron emission scanning in young people shows that the periaqueductal grey matter, pons, and ventral and dorsal portions of the pontine tegmentum are active during bladder filling⁷⁵. Functional magnetic resonance imaging studies in older people suggest that failure of activation in areas of the brain relating to continence, such as the orbitofrontal regions and the insula may lessen the ability to suppress urgency⁷⁶. There is a known association between vascular risk factors and LUTS⁶³, and the presence of white matter hyperintensities (WMH) within periventricular and subcortical regions of the brain is associated with functional and cognitive impairment, an increased incidence of urinary urgency and detrusor overactivity on cystometry and a difficulty in maintaining continence^{77,78}. It has been suggested that changes in the ageing brain, such as the accumulation of white matter hyperintensities, lead to a failure to suppress the physiological sensation of bladder filling, leading to urgency and urgency incontinence⁷⁹. Older adults with vascular dementia are more likely to have LUTS, specifically urgency and urgency incontinence, than those with Alzheimer’s disease⁸⁰. Those with high WMH burden in the frontal lobes are more likely to have incontinence, more severe incontinence, and higher symptom bother⁷⁷. Likewise, the presence of WMH is associated with other geriatric syndromes including cognitive impairment⁸¹ and falls, with data from cross-sectional studies demonstrating that the burden of WMH correlates with frequency of falls⁸², and longitudinal data suggesting that the progression of WMH is associated with an increase in the risk of falls in older people over time⁸³.

THE ROLE OF EXECUTIVE FUNCTION IN GAIT AND CONTINENCE

Executive function (EF) can be defined as “a variety of cognitive processes that use and modify information from many cortical sensory systems in the anterior and posterior brain regions to modulate and produce behaviour”⁸⁴. EF covers a broad range of cognitive tasks, divided by Lezak into “volition, planning, purposive action, and action monitoring”⁸⁵. The areas identified by functional brain imaging as being important in the maintenance of continence and the suppression of urgency are also those involved in handling executive function, and a study in community-dwelling women aged 60 found that those with impaired executive function, tested with 5 separate tests, were more likely to have urgency incontinence than other LUTS⁸⁶.

Likewise, intact executive function is a necessary component of safe walking⁸⁴. Walking and the avoidance of falling are not automatic tasks; they require continuous cognitive input and control^{87,88}. In particular, “successful walking”, defined as the ability to get from point to point without falling or fearing falling through instability, is dependent on executive function⁸⁴. Impairment of executive function has been shown to have a negative effect on gait, with associated increased risk of falls; the InCHIANTI study found that cognitively intact older adults who performed poorly on the trail-marking test, a validated test of executive function, had a lower self-selected gait speed when walking over an obstacle course⁸⁹.

A key aspect of EF is the ability to dual task, that is, to manage simultaneous tasks which divert attention. Diverted attention is the condition in which performing two tasks simultaneously leads to deterioration in performance of one or both tasks, referred to by some as dual task costs⁹⁰. There are three main categories of explanation proposed for this phenomenon; capacity sharing, a model which suggests that the brain has a finite capacity for global function, and if simultaneous tasks exceed this threshold, performance declines⁹¹; the bottleneck (or task-switching) model which suggests that individual brain areas can only perform one function at a time, so if competing tasks require the same pathway, a bottleneck occurs, slowing processing and the cross-talk model, which suggests that simultaneous tasks are more difficult if they both require similar sensory input⁹⁰. EF is felt to be predominantly a function of the frontal and prefrontal cortices, those areas involved in the maintenance of continence, but other areas of the brain, including the limbic system and parietal lobes are also involved⁸⁴.

There is evidence from continent, middle-aged women that the sensation of a strong desire to void induces changes in gait, including an increase in stride-time variability and reduction in gait velocity³³, changes which are associated with increased falls risk⁹². Performance of a distraction task while attempting voluntary contraction of the urethral and anal sphincters causes a reduction in the strength of the contraction as measured with sphincter electromyography⁹³. A Japanese study found that, in a cohort of women attending a clinic with menopausal symptoms, those with urgency incontinence had greater reaction times, as measured with a ruler-drop test, than women with other subtypes of incontinence⁹⁴, although no conclusion can be drawn on the direction of effect for this association. Even in healthy, continent young people a strong desire to void has deleterious effects on cognition; a small study using the strong desire to void as a model for pain found that healthy volunteers who were asked to drink fluid until they experienced an “intense urge” to void had significantly worse cognitive performance on tests of detection, visual attention, and working memory compared to both baseline and post-voiding states⁹⁵. In older adults with incontinence, a multicomponent intervention comprising pelvic floor training and video game dancing improved dual-task performance in women aged over 65, with subgroup analysis suggesting that the largest improvement on the n-back test, a measure of dual task ability, was in women with incontinence⁹⁶. In a focus group based study of adults aged 70 years and older, the women involved identified the desire to void as a source of distraction that may affect the performance of other tasks⁹⁷.

It can be hypothesised that the sensation of urinary urgency, or the strong desire to void, acts as a source of diverted attention, and that other simultaneous cognitive tasks require dual tasking (at cognitive cost) to complete. Given that dual tasking is a well-recognised cause of gait changes and increased falls risk, it may be that this is one underlying mechanism linking urinary urgency and falls. If urinary urgency is a source of attentional demand, then given that that dual-task training can

reduce falls risk under dual task conditions⁹⁸⁻¹⁰¹ there is the potential for novel interventions to address the urgency and linked falls risk.

A MODEL FOR FUTURE RESEARCH

The oft-quoted explanation that urinary urgency causes falls due to rushing to the toilet is an intuitive response, but not supported by the evidence. The temporal relationship between falls and LUTS is far from well-established^{25,32}, and the response to a desire to void appears to be to slow down, not speed up³³. The “rushing” hypothesis has not been subjected to systematic examination and should, therefore, no longer be accepted until there is evidence to support this.

There is good evidence that older adults with LUTS are more likely to have lower levels of physical activity, and that people with lower levels of physical activity are more likely to have LUTS, but the direction of this association is unclear. Further research, both epidemiological and interventional, should be considered to delineate and investigate the role of activity restriction on LUTS and the maintenance of continence in older adults and the role of LUTS and incontinence on activity restriction.

Potential relationships and linkages between LUTS and falls risk factors are outlined in figure 2. There are currently many unexplained steps in the relationship between falls and LUTS with a rich research agenda to be explored. The potential role of urgency as a source of attentional demand can be investigated by comparing the performance of various tasks, such as walking or cognitive tests under conditions of no distraction, urinary urgency, and a validated source of diverted attention such as the n-back test. There is a need for further high-quality epidemiological work to disentangle the influences of frailty, multimorbidity and other factors in LUTS and falls, and it remains unknown if current treatments for OAB, including conservative and pharmacological treatments, either reduce or increase the risk of falling in older people¹⁰². The development of functional brain imaging may provide increasingly sophisticated understanding of the central control of continence, and as our understanding of the underlying physiology of OAB and other LUTS improves, further avenues of study will become apparent. It is clear that WMH are an important common cause of both LUTS and falls, and as such including MRI in studies examining the links between falls and LUTS would be valuable to exclude this factor from analysis.

Despite having a high prevalence of both falls and LUTS, frail and institutionalised older adults are often excluded from research. It is important that researchers consider the needs this population and include them, where possible, in future research.

As Donald Rumsfeld famously told us, the investigation of unknown unknowns is challenging. Clearer data regarding the circumstances of falls in those with LUTS, preferably through robust, prospective data collection, would allow the identification of other potential avenues for study.

There is a rich and unexplored research agenda, and we encourage the development of research projects and programmes to explore these potential links. Does gait change in response to urgency? What changes are observable? Is there a reliable way to induce “urgency” in a gait lab? What underlies this change? Are cognitive factors involved? Are there observable changes to the pelvic floor when walking with and without urgency? Can we ameliorate the associated risk of falls by

treating urgency? This is by its nature a multidisciplinary problem and the opportunities for cross-professional collaboration are large.

CONCLUSIONS

There is a clear association between LUTS, in particular nocturia, urinary urgency and urgency incontinence, and falls in older adults, with significant associated morbidity, mortality, and healthcare resource use. It is not clear, however, to what extent this relationship is due to falls and LUTS having a common cause, and how much is due to factors such as dual tasking, activity restriction, and other, as yet unrecognised, mechanisms. There is little evidence supporting the commonly-held belief that this relationship can be explained through rushing to get to a toilet, and there is evidence to support there being at least an element of “common cause”; that both LUTS and UI are common in later life and very common among the frail. Moving forward using the model we outline will facilitate a greater sophistication in understanding of how these factors are linked and identification of novel interventions to test.

ACKNOWLEDGEMENTS

We are grateful to the attendees of the initial consensus meeting, Dr Derek Griffiths, Mr Justin Lewicke, Dr Danny Rafferty and Dr Sylvie Nadeau, and to the International Continence Society for funding the meeting.

References

1. Irwin DE, Milsom I, Hunskaar S, et al. Population-based survey of urinary incontinence, overactive bladder, and other lower urinary tract symptoms in five countries: results of the EPIC study. *Eur Urol*. Dec 2006;50(6):1306-1314; discussion 1314-1305.
2. Coyne KS, Sexton CC, Thompson CL, et al. The prevalence of lower urinary tract symptoms (LUTS) in the USA, the UK and Sweden: results from the Epidemiology of LUTS (EpiLUTS) study. *BJU Int*. Aug 2009;104(3):352-360.
3. Elstad EA, Taubenberger SP, Botelho EM, Tennstedt SL. Beyond incontinence: the stigma of other urinary symptoms. *J Adv Nurs*. Nov 2010;66(11):2460-2470.
4. Teunissen D, van Weel C, Lagro-Janssen T. Urinary incontinence in older people living in the community: examining help-seeking behaviour. *Br J Gen Pract*. Oct 2005;55(519):776-782.
5. Horrocks S, Somerset M, Stoddart H, Peters TJ. What prevents older people from seeking treatment for urinary incontinence? A qualitative exploration of barriers to the use of community continence services. *Fam Pract*. Dec 2004;21(6):689-696.
6. Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn*. 2002;21(2):167-178.
7. Hester AL, Wei F. Falls in the community: state of the science. *Clin Interv Aging*. 2013;8:675-679.
8. Falls. World Health Organization; 2016.
9. Davis JC, Robertson MC, Ashe MC, Liu-Ambrose T, Khan KM, Marra CA. International comparison of cost of falls in older adults living in the community: a systematic review. *Osteoporos Int*. Aug 2010;21(8):1295-1306.
10. Stel VS, Smit JH, Pluijm SM, Lips P. Consequences of falling in older men and women and risk factors for health service use and functional decline. *Age Ageing*. Jan 2004;33(1):58-65.
11. Collerton J, Kingston A, Bond J, et al. The personal and health service impact of falls in 85 year olds: cross-sectional findings from the Newcastle 85+ cohort study. *PLoS One*. 2012;7(3):e33078.
12. Burns ER, Stevens JA, Lee R. The direct costs of fatal and non-fatal falls among older adults - United States. *J Safety Res*. Sep 2016;58:99-103.
13. Scott V, Wagar L, Elliot S. Falls & Related Injuries among Older Canadians: Fall-related Hospitalizations & Intervention Initiatives. In: Public Health Agency of Canada DoAaS, ed. Victoria, BC.2010.
14. Brown JS, Vittinghoff E, Wyman JF, et al. Urinary incontinence: does it increase risk for falls and fractures? Study of Osteoporotic Fractures Research Group. *J Am Geriatr Soc*. Jul 2000;48(7):721-725.
15. Chiarelli PE, Mackenzie LA, Osmotherly PG. Urinary incontinence is associated with an increase in falls: a systematic review. *Aust J Physiother*. 2009;55(2):89-95.
16. Damian J, Pastor-Barriuso R, Valderrama-Gama E, de Pedro-Cuesta J. Factors associated with falls among older adults living in institutions. *BMC Geriatr*. 2013;13:6.
17. Foley AL, Loharuka S, Barrett JA, et al. Association between the Geriatric Giants of urinary incontinence and falls in older people using data from the Leicestershire MRC Incontinence Study. *Age Ageing*. Jan 2012;41(1):35-40.
18. Galizia G, Langellotto A, Cacciatore F, et al. Association between nocturia and falls-related long-term mortality risk in the elderly. *J Am Med Dir Assoc*. Sep 2012;13(7):640-644.
19. Health Quality O. Behavioural interventions for urinary incontinence in community-dwelling seniors: an evidence-based analysis. *Ont Health Technol Assess Ser*. 2008;8(3):1-52.
20. Hedman AM, Fonad E, Sandmark H. Older people living at home: associations between falls and health complaints in men and women. *J Clin Nurs*. Oct 2013;22(19-20):2945-2952.

21. Lee CY, Chen LK, Lo YK, et al. Urinary incontinence: an under-recognized risk factor for falls among elderly dementia patients. *Neurourol Urodyn*. Sep 2011;30(7):1286-1290.
22. Brown JS, Vittinghoff E, Wyman JF. Weekly urge urinary incontinence was associated with increased risk for falls and non-spinal fractures in older women. *Evidence-Based Medicine*. 2001;6(2):59.
23. Wagner TH, Hu T, Bentkover J, et al. Health-related consequences of overactive bladder. *American Journal of Managed Care*. 2002;8(19):S598-607.
24. Noguchi N, Chan L, Cumming RG, et al. Lower Urinary Tract Symptoms and Incident Falls in Community Dwelling Older Men: The Concord Health and Ageing in Men Project. *J Urol*. Jun 24 2016.
25. Sakushima K, Yamazaki S, Fukuma S, et al. Influence of urinary urgency and other urinary disturbances on falls in Parkinson's disease. *J Neurol Sci*. Jan 15 2016;360:153-157.
26. Noguchi N, Chan L, Cumming RG, Blyth FM, Naganathan V. A systematic review of the association between lower urinary tract symptoms and falls, injuries, and fractures in community-dwelling older men. *Aging Male*. Apr 12 2016:1-7.
27. Pahwa AK, Andy UU, Newman DK, Stambakio H, Schmitz KH, Arya LA. Nocturnal Enuresis as a Risk Factor for Falls in Older Community Dwelling Women with Urinary Incontinence. *J Urol*. May 2016;195(5):1512-1516.
28. Godmaire GC, Grenier S, Tannenbaum C. An independent association between urinary incontinence and falls in chronic benzodiazepine users. *J Am Geriatr Soc*. May 2015;63(5):1035-1037.
29. Morris V, Hunter K, Wagg A. Falls and urinary incontinence: a link ripe for intervention? *GM: Midlife & Beyond*. 2011;41(6):333-336.
30. Morris V, Wagg A. Lower urinary tract symptoms, incontinence and falls in elderly people: time for an intervention study. *Int J Clin Pract*. Feb 2007;61(2):320-323.
31. Wagner TH, Hu TW. Economic costs of urinary incontinence in 1995. *Urology*. Mar 1998;51(3):355-361.
32. Edwards R, Hunter K, Wagg A. Lower urinary tract symptoms and falls in older women: A case control study. *Maturitas*. Dec 24 2014.
33. Booth J, Paul L, Rafferty D, Macinnes C. The relationship between urinary bladder control and gait in women. *Neurourol Urodyn*. Jan 2013;32(1):43-47.
34. Jefferis BJ, Iliffe S, Kendrick D, et al. How are falls and fear of falling associated with objectively measured physical activity in a cohort of community-dwelling older men? *BMC Geriatr*. Oct 27 2014;14:114.
35. Murphy SL, Williams CS, Gill TM. Characteristics associated with fear of falling and activity restriction in community-living older persons. *J Am Geriatr Soc*. Mar 2002;50(3):516-520.
36. Fletcher PC, Hirdes JP. Restriction in activity associated with fear of falling among community-based seniors using home care services. *Age Ageing*. May 2004;33(3):273-279.
37. Delbaere K, Crombez G, Vanderstraeten G, Willems T, Cambier D. Fear-related avoidance of activities, falls and physical frailty. A prospective community-based cohort study. *Age Ageing*. Jul 2004;33(4):368-373.
38. Brown WJ, Miller YD. Too wet to exercise? Leaking urine as a barrier to physical activity in women. *J Sci Med Sport*. Dec 2001;4(4):373-378.
39. Nelson PR, Irish KR, Cleary KK. A Preliminary Study on Balance Performance and Fall Status in Older Women With Urinary Incontinence. *Journal of Women's Health Physical Therapy*. September/December 2015 2015;39(3):102-108.
40. Reigota RB, Pedro AO, de Souza Santos Machado V, Costa-Paiva L, Pinto-Neto AM. Prevalence of urinary incontinence and its association with multimorbidity in women aged 50 years or older: A population-based study. *Neurourol Urodyn*. Jan 2016;35(1):62-68.

41. McGrother CW, Donaldson MM, Hayward T, et al. Urinary storage symptoms and comorbidities: a prospective population cohort study in middle-aged and older women. *Age Ageing*. Jan 2006;35(1):16-24.
42. Danforth KN, Shah AD, Townsend MK, et al. Physical activity and urinary incontinence among healthy, older women. *Obstet Gynecol*. Mar 2007;109(3):721-727.
43. Fritel X, Lachal L, Cassou B, Fauconnier A, Dargent-Molina P. Mobility impairment is associated with urge but not stress urinary incontinence in community-dwelling older women: results from the Ossebo study. *BJOG*. Nov 2013;120(12):1566-1572.
44. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing*. Sep 2006;35 Suppl 2:ii37-ii41.
45. Jirovec MM. The impact of daily exercise on the mobility, balance and urine control of cognitively impaired nursing home residents. *International journal of nursing studies*. 1991;28(2):145-151.
46. Kim H, Yoshida H, Suzuki T. The effects of multidimensional exercise treatment on community-dwelling elderly Japanese women with stress, urge, and mixed urinary incontinence: a randomized controlled trial. *Int J Nurs Stud*. Oct 2011;48(10):1165-1172.
47. van Houten P, Achterberg W, Ribbe M. Urinary incontinence in disabled elderly women: a randomized clinical trial on the effect of training mobility and toileting skills to achieve independent toileting. *Gerontology*. 2007;53(4):205-210.
48. Gillespie LD, Robertson MC, Gillespie WJ, et al. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev*. 2009(2):CD007146.
49. Cameron ID, Gillespie LD, Robertson MC, et al. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst Rev*. 2012;12:CD005465.
50. Burton E, Cavalheri V, Adams R, et al. Effectiveness of exercise programs to reduce falls in older people with dementia living in the community: a systematic review and meta-analysis. *Clin Interv Aging*. 2015;10:421-434.
51. Davis JC, Robertson MC, Ashe MC, Liu-Ambrose T, Khan KM, Marra CA. Does a home-based strength and balance programme in people aged > or =80 years provide the best value for money to prevent falls? A systematic review of economic evaluations of falls prevention interventions. *Br J Sports Med*. Feb 2010;44(2):80-89.
52. Tinetti ME, Inouye SK, Gill TM, Doucette JT. Shared risk factors for falls, incontinence, and functional dependence: Unifying the approach to geriatric syndromes. *Journal of the American Medical Association*. 1995;273(17):1348-1353.
53. Nowak A, Hubbard RE. Falls and frailty: lessons from complex systems. *J R Soc Med*. Mar 2009;102(3):98-102.
54. Ensrud KE, Ewing SK, Taylor BC, et al. Frailty and risk of falls, fracture, and mortality in older women: the study of osteoporotic fractures. *J Gerontol A Biol Sci Med Sci*. Jul 2007;62(7):744-751.
55. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. Mar 2001;56(3):M146-156.
56. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci*. Jul 2007;62(7):722-727.
57. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ*. Aug 30 2005;173(5):489-495.
58. Rolfson DB, Majumdar SR, Tsuyuki RT, Tahir A, Rockwood K. Validity and reliability of the Edmonton Frail Scale. *Age Ageing*. Sep 2006;35(5):526-529.
59. Kim H, Yoshida H, Hu X, et al. Association between self-reported urinary incontinence and musculoskeletal conditions in community-dwelling elderly women: A cross-sectional study. *Neurology and Urodynamics*. 2014.

60. Gosch M, Talasz H, Nicholas JA, Kammerlander C, Lechleitner M. Urinary incontinence and poor functional status in fragility fracture patients: an underrecognized and underappreciated association. *Arch Orthop Trauma Surg.* Jan 2015;135(1):59-67.
61. Thom DH, Haan MN, Van Den Eeden SK. Medically recognized urinary incontinence and risks of hospitalization, nursing home admission and mortality. *Age Ageing.* Sep 1997;26(5):367-374.
62. Herr M, Robine JM, Pinot J, Arvieu JJ, Ankri J. Polypharmacy and frailty: prevalence, relationship, and impact on mortality in a French sample of 2350 old people. *Pharmacoepidemiol Drug Saf.* Jun 2015;24(6):637-646.
63. Ponholzer A, Temml C, Wehrberger C, Marszalek M, Madersbacher S. The association between vascular risk factors and lower urinary tract symptoms in both sexes. *Eur Urol.* Sep 2006;50(3):581-586.
64. Franceschi C, Campisi J. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J Gerontol A Biol Sci Med Sci.* Jun 2014;69 Suppl 1:S4-9.
65. Hashimoto M, Hashimoto K, Ando F, Kimura Y, Nagase K, Arai K. Prescription rate of medications potentially contributing to lower urinary tract symptoms and detection of adverse reactions by prescription sequence symmetry analysis. *J Pharm Health Care Sci.* 2015;1:7.
66. Kashyap M, Tu le M, Tannenbaum C. Prevalence of commonly prescribed medications potentially contributing to urinary symptoms in a cohort of older patients seeking care for incontinence. *BMC Geriatr.* Jun 10 2013;13:57.
67. Devore EE, Townsend MK, Resnick NM, Grodstein F. The epidemiology of urinary incontinence in women with type 2 diabetes. *J Urol.* Nov 2012;188(5):1816-1821.
68. Mayne D, Stout NR, Aspray TJ. Diabetes, falls and fractures. *Age Ageing.* Sep 2010;39(5):522-525.
69. Tsujimura A, Yamamoto Y, Sakoda S, et al. Finger taps and constipation are closely related to symptoms of overactive bladder in male patients with Parkinson's disease. *International Journal of Urology.* 2014;21(1):69-73.
70. Wood BH, Bilclough JA, Bowron A, Walker RW. Incidence and prediction of falls in Parkinson's disease: a prospective multidisciplinary study. *J Neurol Neurosurg Psychiatry.* Jun 2002;72(6):721-725.
71. Brown JS, Vittinghoff E, Wyman JF, et al. Urinary incontinence: does it increase risk for falls and fractures? *Journal of the American Geriatrics Society.* 2000;48(7):721-725.
72. Zarowitz BJ, Allen C, O'Shea T, Tangalos E, Berner T, Ouslander JG. Clinical Burden and Nonpharmacologic Management of Nursing Facility Residents with Overactive Bladder and/or Urinary Incontinence. *Consult Pharm.* Sep 2015;30(9):533-542.
73. Wu HY. Achieving urinary continence in children. *Nat Rev Urol.* Jul 2010;7(7):371-377.
74. Fowler CJ, Griffiths DJ. A decade of functional brain imaging applied to bladder control. *Journal of Urology.* 2011;186(2):615-616.
75. Matsuura S, Kakizaki H, Mitsui T, Shiga T, Tamaki N, Koyanagi T. Human brain region response to distention or cold stimulation of the bladder: a positron emission tomography study. *J Urol.* Nov 2002;168(5):2035-2039.
76. Griffiths D, Tadic SD, Schaefer W, Resnick NM. Cerebral control of the bladder in normal and urge-incontinent women. *Neuroimage.* Aug 1 2007;37(1):1-7.
77. Kuchel GA, Moscufo N, Guttmann CR, et al. Localization of brain white matter hyperintensities and urinary incontinence in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci.* Aug 2009;64(8):902-909.
78. Kuo HK, Lipsitz LA. Cerebral white matter changes and geriatric syndromes: is there a link? *J Gerontol A Biol Sci Med Sci.* Aug 2004;59(8):818-826.

79. Resnick N, Tadic S, Clarkson B, Murrin A, Griffiths D. Brain mechanisms underlying behavioral therapy for urgency incontinence in older women. *43rd Annual Meeting of the International Continence Society, ICS 2013 Barcelona Spain*. 2013;32(6):795-796.
80. Takahashi O, Sakakibara R, Panicker J, et al. White matter lesions or Alzheimer's disease: Which contributes more to overactive bladder and incontinence in elderly adults with dementia? *Journal of the American Geriatrics Society*. 2012;60(12):2370-2371.
81. Prins ND, Scheltens P. White matter hyperintensities, cognitive impairment and dementia: an update. *Nat Rev Neurol*. Mar 2015;11(3):157-165.
82. Baezner H, Blahak C, Poggesi A, et al. Association of gait and balance disorders with age-related white matter changes: the LADIS study. *Neurology*. Mar 18 2008;70(12):935-942.
83. Callisaya ML, Beare R, Phan T, et al. Progression of white matter hyperintensities of presumed vascular origin increases the risk of falls in older people. *J Gerontol A Biol Sci Med Sci*. Mar 2015;70(3):360-366.
84. Yogev-Seligmann G, Hausdorff JM, Giladi N. The role of executive function and attention in gait. *Mov Disord*. Feb 15 2008;23(3):329-342; quiz 472.
85. Lezak MD. *Neuropsychological assessment*. 5th ed. Oxford ; New York: Oxford University Press; 2012.
86. Lussier M, Renaud M, Chiva-Razavi S, Bherer L, Dumoulin C. Are stress and mixed urinary incontinence associated with impaired executive control in community-dwelling older women? *J Clin Exp Neuropsychol*. 2013;35(5):445-454.
87. Verghese J, Kuslansky G, Holtzer R, et al. Walking while talking: effect of task prioritization in the elderly. *Arch Phys Med Rehabil*. Jan 2007;88(1):50-53.
88. Shumway-Cook A, Woollacott M. Attentional demands and postural control: the effect of sensory context. *J Gerontol A Biol Sci Med Sci*. Jan 2000;55(1):M10-16.
89. Ble A, Volpato S, Zuliani G, et al. Executive function correlates with walking speed in older persons: the InCHIANTI study. *J Am Geriatr Soc*. Mar 2005;53(3):410-415.
90. Pashler H. Dual-task interference in simple tasks: data and theory. *Psychol Bull*. Sep 1994;116(2):220-244.
91. Tombu M, Jolicoeur P. A central capacity sharing model of dual-task performance. *J Exp Psychol Hum Percept Perform*. Feb 2003;29(1):3-18.
92. Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil*. Aug 2001;82(8):1050-1056.
93. Thubert T, Deffieux X, Jousse M, Guinet-Lacoste A, Ismael SS, Amarenco G. Influence of a distraction task on pelvic floor muscle contraction. *Neurourol Urodyn*. Feb 2015;34(2):139-143.
94. Terauchi M, Hirose A, Akiyoshi M, Owa Y, Kato K, Kubota T. Prevalence and predictors of storage lower urinary tract symptoms in perimenopausal and postmenopausal women attending a menopause clinic. *Menopause*. Oct 2015;22(10):1084-1090.
95. Lewis MS, Snyder PJ, Pietrzak RH, Darby D, Feldman RA, Maruff P. The effect of acute increase in urge to void on cognitive function in healthy adults. *Neurourol Urodyn*. Jan 2011;30(1):183-187.
96. Fraser SA, Elliott V, de Bruin ED, Bherer L, Dumoulin C. The Effects of Combining Videogame Dancing and Pelvic Floor Training to Improve Dual-Task Gait and Cognition in Women with Mixed-Urinary Incontinence. *Games Health J*. Jun 2014;3(3):172-178.
97. Muhaidat J, Skelton DA, Kerr A, Evans JJ, Ballinger C. Older adults' experiences and perceptions of dual tasking. *The British Journal of Occupational Therapy*. 2010 2010;73(9):405-412.
98. Li KZ, Roudaia E, Lussier M, Bherer L, Leroux A, McKinley PA. Benefits of cognitive dual-task training on balance performance in healthy older adults. *J Gerontol A Biol Sci Med Sci*. Dec 2010;65(12):1344-1352.

99. Pichierri G, Coppe A, Lorenzetti S, Murer K, de Bruin ED. The effect of a cognitive-motor intervention on voluntary step execution under single and dual task conditions in older adults: a randomized controlled pilot study. *Clin Interv Aging*. 2012;7:175-184.
100. Schwenk M, Zieschang T, Oster P, Hauer K. Dual-task performances can be improved in patients with dementia: a randomized controlled trial. *Neurology*. Jun 15 2010;74(24):1961-1968.
101. Silsupadol P, Lugade V, Shumway-Cook A, et al. Training-related changes in dual-task walking performance of elderly persons with balance impairment: a double-blind, randomized controlled trial. *Gait Posture*. Jun 2009;29(4):634-639.
102. Gomes T, Juurlink DN, Ho JM, Schneeweiss S, Mamdani MM. Risk of serious falls associated with oxybutynin and tolterodine: a population based study. *J Urol*. Oct 2011;186(4):1340-1344.

Figure 1: Common causes of falls and LUTS in older adults




Figure 2: Potential (dashed line), known (solid line), and suggested but likely incorrect (dotted line) links between urgency/LUTS and falls

